009068 ARH-LD-132 Informal Report

-201

REFERENCE COPY
MUT A CONCRETED APR 2 8 1976
RECEIVED - ARRICO
DOCUMENT CONCRETE
354 Federal, Richland

ENTERED

Geology of the 241-C Tank Farm

April 1976

W. H. Price

K. R. Fecht



Environmental Engineering Section Research Department Research and Engineering Division

Prepared for the U.S. Energy Research and Development Administration Under Contract E(45-1)-2130

Atlantic Richfield Hanford Company Richland, Washington 99352



GEOLOGY OF THE 241-C TANK FARM

bу

W. H. Price K. R. Fecht

Environmental Engineering Section Research Department Research and Engineering Division

April 1976

ATLANTIC RICHFIELD HANFORD COMPANY RICHLAND, WASHINGTON 99352

TABLE OF CONTENTS

																					Page
INTRO	ODUC.	TION.	•	•	•	•	•	•	•	•	•	•		•							1
PROCE	EDUR	ES .	• •		•	•	•	•	•	•		•	•		•	•					2
GENE	RAL I	ZED GE	OLOGY	· .	•	•		•	•	•				•			•			•	3
	COL	JMBIA	RIVER	BAS	SALT	GF	OUP		•	•		•		•	•		•	•	•		6
	RING	GOLD F	ORMAT	ION			•				•	•	•	•	•	•	•	•			6
	GLA	CIOFLU	VIAL	DEPO	SIT	s.		•	•				•		•	•		•			6
	CLAS	STIC D	IKES.	•					•	•			•	•							8
	BACK	FILL	MATER	IAL			•		•		•			•					·	•	8
	WATE	R TAB	LE .		•	•	•	•		•	•				_			•	•	•	8
GLOSS	SARY			•						•						•	•	•	•	•	9
SELEC	TED	REFER	ENCES	•	•				•				•		•	•	•	•	•	•	11
											•	•	•	•	•	•	•	•	•	•	11
							LIS	T	OF	TABI	LES										
Table																					
I	•	TANK	FARM	GEOL	ngv	י חר	CHM	E NI	TC /	11/4	t. A.		• •		•						Page
II III			FARM TANK	N CAP	KIM G	たいし	.UGY	M/	166											•	1 2
IV		GEACT	CAL GR	IAL	LII	HUL	DG L	FS	REN	IF A 7	TH 2	M1_	C 7	TARIL	/ EA	DM		1 AJ(OR •		7
		11610	CAL GR 241-C	CATM	217	E A	עומו	CAL	CH	IM (:ARB	ONA	TE.	VAL.	.UES	. F0	R •				8
																				•	Ū
							LIS	TC)F F	IGU	IRES										
Figure	<u> </u>																				Page
1		STEPS	OUTL	ININ	G TI	HE	PREI	PAR	ATI	ON	OF	TAN	K								. 494
2		FARM GENER	GEOLO: ALIZE	GY M. DST	APS RAT:	IGR	APH		_					•	•	•	•	•	•		4
		200 A	REA T	ANK	FAR	YS	•	•	•	•	•	•	•		•		•				5

GEOLOGY OF THE 241-C TANK FARM

INTRODUCTION

A series of maps have been compiled to document the structure and stratigraphy of the sediments underlying the high-level radioactive waste storage tank farms located within the Energy Research and Development Administration Hanford Reservation. The primary purpose of these maps is to provide basic geologic information to be utilized to evaluate the impact of suspected and confirmed tank leaks. For convenience of usage map sets for each tank farm have been published in separate document packets (see Table I). The contents of this packet (see Table II) contain maps compiled only for the 241-C Tank Farm.

TABLE I

TANK FARM GEOLOGY DOCUMENTS AVAILABLE
AS OF APRIL, 1976*

Title	Document Number
Geology of the 241-A Tank Farm	ARH-LD-127
Geology of the 241-AX Tank Farm	ARH-LD-128
Geology of the 241-B Tank Farm	ARH-LD-129
Geology of the 241-BX Tank Farm	ARH-LD-130
Geology of the 241-BY Tank Farm	ARH-LD-131
Geology of the 241-C Tank Farm	ARH-LD-132
Geology of the 241-S Tank Farm	ARH-LD-133
Geology of the 241-SX Tank Farm	ARH-LD-134
Geology of the 241-T Tank Farm	ARH-LD-135
Geology of the 241-TX Tank Farm	ARH-LD-136
Geology of the 241-TY Tank Farm	ARH-LD-137
Geology of the 241-U Tank Farm	ARH-LD-138
Generalized Geology of the	
241-31 lank farm	ARH-LD-139
*Additional documents will be seen	-4.1

^{*}Additional documents will be completed as new tank farms are built and well monitoring networks installed.

TABLE II

241-C TANK FARM GEOLOGY MAPS

Title	Drawing Number
241-C Tank Farm Geologic Map Legend and Plot Plan	H-2-38983
241-C Tank Farm Geologic Characterization Cross Section A-A'	H-2-70496
241-C Tank Farm Geologic Characterization Cross Section B-B'	H-2-70497
241-C Tank Farm Geologic Characterization Cross Section C-C'	H-2-70498
241-C Tank Farm Geologic Characterization Cross Section D-D'	H-2-70499
241-C Tank Farm Geologic Characterization Cross Section E-E'	H-2-70500
241-C Tank Farm Geologic Characterization Cross Section F-F'	H-2-70501
241-C Tank Farm Geologic Characterization Cross Section G-G'	H-2-70502
241-C Tank Farm Geologic Characterization Cross Section H-H'	H-2-70504
241-C Tank Farm Geologic Characterization Cross Section I-I'	H-2-70503
241-C Tank Farm Geologic Characterization Base of Backfill	H-2-70495

PROCEDURES

During the drilling of 26 dry wells and 2 water wells in and around the 241-C Tank Farm, sediment samples were collected from one to 5-foot depth intervals. Information utilized to prepare this series of maps was obtained by the analysis of these samples, numbering approximately 600.

Each sediment sample was quantitatively analyzed according to grain size and ${\rm CaCO}_3$ content. Size analysis was carried out utilizing a nest of 9 sieves selected for coincidence with the Wentworth (1922) grain size nomenclature (see H-2-38983). The ${\rm CaCO}_3$ content of each sample was determined utilizing a semiquantitative ${\rm CO}_2$ displacement method (Horwitz, 1970). Size and ${\rm CaCO}_3$ data was input into the Rocksan Computer Program (Parr, 1974) which categorized each sediment sample into 1 of 19 classes (classification scheme modified after Folk, 1968; see H-2-38983). After analysis, each sample was visually examined to aid in further characterization. Each sample was subsequently stored in the Hanford Well Library for future reference.

For convenience of usage, the geologic maps were prepared at the same scale (1"=16') as drawing H-2-36941 (Wells in 241-C Farm As-built). Steps outlining the preparation of the maps are listed in Figure 1.

GENERALIZED GEOLOGY

Included within this section is a brief discussion of the geology underlying the 241-C Tank Farm. The stratigraphic descriptions included, along with the Glossary (see page 9), are designed only to provide sufficient information to permit a general understanding of the Tank Farm maps presented. For a more detailed discussion of the regional geologic setting of the 241-C Tank Farm, the reader is referred to articles listed in the Selected References (see page 11).

The 241-C Tank Farm is underlain by three major stratigraphic units (see Figure 2); (1) basalt of the Columbia River Group which forms the bedrock beneath the area; (2) semiconsolidated sediments of the Ringold Formation which directly overlie the bedrock; and (3) unconsolidated sand, silt, and gravel, collectively termed glaciofluvial sediments. A more detailed description of the character of these units underlying the Tank Farm follows.




FIGURE 1

STEPS OUTLINING THE PREPARATION OF TANK FARM GEOLOGY MAPS

	To	•													
ERA	PERIOD	ЕРОСН	YEARS B. P.	1	TRATIGRAPHIC IAME AND/OR UNIT	LITHOLOGY DESCRIPTION									
		MODERN	30		BACKFILL	VERY POORLY SORTED GRAVEL, SAND & SILT									
	QUATERNARY	PLEISTOCENE	33	GI	ACIOFLUVIAL SEDIMENTS	FAIRLY WELL SORTED FLUVIAL SAND & SILT WITH SOME GRAVEL									
	QUATE	PLEIST	- 1,000,000 -		EOLIAN SILT	FINE SAND & SILT DERIVED FROM THE UPPER RINGOLD									
		NE		RINGOLD FORMATION	UPPER RINGOLD	WELL SORTED FLUVIAL OR LACUSTRINE SILT & SAND WITH SOME CALCAREOUS LAYERS									
CENOZOIC		PLIOCENE		RINGOLD F	MIDDLE RINGOLD	FLUVIAL GRAVEL & SAND VARIABLY CEMENTED WITH CALCIUM CARBONATE & SILICA									
	TERTIARY				11,000,000 -		11,000,000 -	· 11,000,000 -	· 11,000,000 -	11,000,000 -		GROUP		ELEPHANT MOUNTAIN MEMBER	DENSE BLACK EXTRUSIVE IGNEOUS ROCK, MICRO VESICULAR, BRICK BAT ENTABULATURE & NO COLUMNADE
		MIOCENE		ER BASALT	RATTLESNAKE RIDGE MEMBER	TUFFACEOUS SANDSTONE									
		Ž		COLUMBIA RIVER	POMONA MEM BER	DENSE BLACK EXTRUSIVE IGNEOUS ROCK, SCATTERED OLIVINE PHENOCRYSTS, UPPER & SOMETIMES BASAL ENTABLATURE WELL DEVELOPED, FAN JOINTING IN COLUMNADE									

WHP/KRF 1976

FIGURE 2

GENERALIZED STRATIGRAPHIC COLUMN FOR THE 200 AREA TANK FARMS

COLUMBIA RIVER BASALT GROUP

About 20 million years ago a series of fissures opened around the periphery of the subsiding Pasco Basin and large volumes of basaltic lava poured out over the land surface. The highly fluid lava was extruded intermittently from these fissures until approximately 8 million years ago. At the cessation of Columbia River Basalt volcanism, the basin had been filled with more than 12,000 feet of basalt.

The surface of the Columbia River Basalt lies beneath 241-C Tank Farm at an elevation of approximately 330 feet (all elevations based on feet above mean sea level measured at approximate center of Tank Farm). On the 241-C Tank Farm maps, this surface occurs approximately 30 feet below the bottom border of the prepared cross sections.

RINGOLD FORMATION

Following the cessation of Columbia River Basalt volcanism the ancestral Columbia River transported sediments from the surrounding highlands into the Pasco Basin where they accumulated to form the Ringold Formation. Beneath the Hanford Reservation, this formation is up to 1200 feet thick and can generally be divided into three units on the basis of lithology; the clays and silts of the lower Ringold unit; the pebbles and cobbles of the middle Ringold unit; and the silts and fine sands of the upper Ringold unit.

Within the region beneath 241-C Tank Farm, the total thickness of the Ringold Formation is estimated to be 20 feet. Sediments samples were not available to adequately describe the Formation.

GLACIOFLUVIAL DEPOSITS

During the close of the Ice Age, approximately 20,000 years ago, a continental ice sheet covered much of northern Washington. As the ice sheet retreated northward, the breakup of ice dams resulted in catastrophic floods in which large volumes of glacial meltwater were released. During one of these floods, over 500 cubic miles of water is estimated to have poured into the Pasco Basin at a rate of more than 9 cubic miles of water per

hour. Sediments deposited within the basin by such flooding now comprise the glaciofluvial unit. The characteristic variability of sediment size and degree of sorting within this unit can be attributed to changes in water velocity and water level which occurred during the flooding process.

Glaciofluvial deposits are found beneath the 241-C Tank Farm between elevations 330 and 607 feet. The 277-foot thick section of these deposits consists predominently of very coarse to coarse sand with some silt and pebbles. Table III summarizes the grain size and ${\rm CaCO}_3$ values of the glaciofluvial sediments.

TABLE III

TYPICAL GRAIN SIZE AND CALCIUM CARBONATE VALUES
FOR MAJOR GLACIOFLUVIAL LITHOLOGIES
BENEATH 241-C TANK FARM

Lithology	%Pebbles & Cobbles	Very Coarse	Coarse	Sand Medium	<u>Fine</u>	Very	%Silt	**	
Slightly Silty Very Coarse to Coarse Sand to Very Coarse to Coarse Sand	1	20	43	14	7	<u>Fine</u>	<u>& Clay</u> 10	%CaCO ₃	
Slightly Pebbly Very Coarse to Coarse Sand	7	29	31	17	7	4	5	2.0	
Pebbly Very Coarse to Coarse Sand	15	26	33	13	6	3	4	1.6	
Slighty Silty Coarse to Medium Sand to Coarse to Medium Sand	3	14	37	24	9	4	9	1.3	
Slightly Silty Very Coarse to Coarse Sand to Very Coarse to Coarse Sand	2	24	31	16	9	9	9	1.2	
Slightly Silty Medium Sand to Coarse to Medium Sand	3	10	28	33	11	6	9	0.8	
Very Coarse to Coarse Sand	1	28	45	14	5	2	5	0.8	
Slightly Pebbly Very Coarse to Coarse Sand	8	27	31	15	6	4	9	0.7	

CLASTIC DIKES

Throughout the Pasco Basin, clastic dikes are found cross-cutting the Ringold Formation and glaciofluvial sediments. These dikes, which range from a few inches to several feet in width, are known to exist to depths of more than 100 feet below the ground surface. Generally, the dikes are composed of fine silts to coarse sands. The origin of the clastic dikes is still in refute and will not be discussed here (see Selected References). Identification of clastic dikes by drilling is difficult and although some dikes were detected in the 241-C Tank Farm, they could not be mapped.

BACKFILL MATERIAL

In preparation for tank construction, glaciofluvial material was excavated at the 241-C Tank Farm site. This material, consisting predominantly of cobbles, pebbles, and coarse to medium sands with some silts, was subsequently used as backfill from the base of the completed tanks (607 feet) to the ground surface (646 feet). An inherent characteristic of the backfill is its poor sorting. Grain size and CaCO₃ values for the backfill are found in Table IV.

TABLE IV

TYPICAL GRAIN SIZE AND CALCIUM CARBONATE VALUES FOR THE 241-C TANK FARM BACKFILL

	%Pebbles	V	% Sand						
Lithology	& Cobbles	Very Coarse	Coarse	Medium	Fine	Very Fine	%Silt & Clay	%CaCO3	
Gravelly Very Coarse to Medium Sand to Slightly Gravelly Very Coarse to Medium Sand		26	17	10	9	7	14	1.6	

WATER TABLE

The water table beneath the 241-C Tank Farm is located within the glacio-fluvial sediments at an elevation of 403 feet, 204 feet below the base of the tanks. For further information concerning contours on the water table beneath 200 East Area the reader is referred to drawings H-2-38398 (200 East Area Water Table Map) and H-2-38399 (200 East Area Depth to Water Map).

GLOSSARY

- Basalt. Fine-grained, dark-colored, extrusive igneous rock.
- <u>Calcareous</u>. Containing calcium carbonate.
- Caliche. Gravel, sand, or silt cemented by calcium carbonate.
- <u>Cement.</u> Chemically precipitated material occurring in the interstices between particles of gravel, sand, or silt.
- <u>Clastic</u>. A textural term applied to rocks composed of fragmental material derived from pre-existing rocks.
- <u>Clastic dike</u>. A tabular body of clastic material transecting the bedding of a sedimentary formation, representing extraneous material that has invaded the containing formation along a crack.
- <u>Dip.</u> The angle at which a stratum or any planar feature is inclined from the horizontal.
- $\frac{\text{Eolian}}{\text{air.}}$ A formation formed by, or deposited from, the wind or currents of
- Fluvial. Produced by the action of a river or stream.
- <u>Formation</u>. The ordinary unit of geologic mapping consisting of a large and persistent stratum of some one kind of rock.
- Glaciofluvial. Pertaining to streams flowing from glaciers or to the deposits made by such streams.
- Grain. The particles or discrete crystals which comprise a rock or sediment.
- Group. A local or provincial subdivision of a series, based on lithologic features and contains two or more formations.
- Lacustrine. A formation deposited in a lake environment.
- Lava. Fluid rock such as that which issues from a volcano or a fissure in the earth's surface and the same material solidified by cooling.
- <u>Lithology</u>. The description of rocks or sediments on the basis of such characteristics as color, minerologic composition and grain size.
- <u>Sediment</u>. Descriptive term for gravel, sand, and silt transported from their sources and deposited by air, water, or ice.
- <u>Sieve</u>. A utensil having many small perforated openings, used to separate fine particles from coarser ones.

Siliceous. Containing silica.

Silt. Fine grained material between sand and clay in size.

Sorting. The grain size range of the sediments.

Stratigraphy. The part of descriptive geology of an area that pertains to the discrimination, character, thickness, sequence, age and correlation of the sediments and rocks of the area.

Subaerial. Formed, existing, or taking place on the land surface.

<u>Unconformity</u>. A surface of erosion or nondeposition that separates younger strata from older strata.

Water table. The upper surface of a zone of saturation except where that surface is formed by an impermeable body.

Winnowing. Separation of fine particles from coarser ones by wind action.

SELECTED REFERENCES

- J. A. Alwin, <u>Clastic Dikes of Touchet Beds</u>, <u>Southeastern Washington</u>, Washington State University Masters Thesis (1970).
- V. R. Baker, <u>Paleohydrology and Sedimentology of Lake Missoula Flooding</u> in <u>Eastern Washington</u>, Geological Society of America Special Paper 144, (1973).
- 3. J. H. Bretz, <u>Washington's Channeled Scabland</u>, Washington Division of Mines and Geology, Bull. 45, (1959).
- 4. D. J. Brown, <u>Subsurface Geology of the Hanford Separations Areas</u>, HW-61780 (1959).
- 5. D. J. Brown, An Eolian Deposit Beneath 200 West Area, HW-67549 (1960).
- 6. D. J. Brown, and R. E. Brown, <u>Touchet Clastic Dikes in the Ringold Formation</u>, HW-SA-2851 (1962).
- 7. D. J. Brown, G. T. Lobdell, and G. E. Neff, <u>Hydrology and Engineering Geology of the Columbia Basin</u>, Geological Society of America, Cordilleran Section, 72nd Annual Meeting, Field Guide No. 3 (1976).
- 8. R. L. Folk, "Petrology of Sedimentary Rocks," University of Texas Press (1968).
- 9. R. Fryxell, and E. F. Cook, <u>A Field Guide to the Loess Deposits and Channeled Scablands of the Palouse Area, Eastern Washington</u>, Laboratory of Anthropology Report 27, Pullman, Washington (1964).
- 10. E. H. Gilmour, and D. Stradling, <u>Proceedings of the Second Columbia River Basalt Symposium</u>, EWSC Press, Cheney, Washington (1969).
- 11. E. Gustafson, <u>A Revised Chronology for Vertebrate Fossil Faunas of Eastern Washington</u>, Geological Society of America, Cordilleran Section, 72nd Annual Meeting (1976).
- 12. W. Horwitz, Official Methods of Analysis of the Association of Official Analytical Chemists, 11th Edition, Association of Official Analytical Chemists, p. 139, (1970).
- 13. R. K. Ledgerwood, D. J. Brown, C. W. Meyers, and A. C. Waters, <u>Identification of Yakima Basalt Flows in the Pasco Basin</u>, ARH-27-68 (1973).
- 14. R. L. Lupher, <u>Clastic Dikes of the Columbia Basin Region</u>, <u>Washington and Idaho</u>, Geological Society of America Bull., V55, 1431-1462 (1944).

- 15. M. W. McGoniea, Deformation of the Ringold Formation, HW-36373 (1955).
- B. McKee, <u>Cascadia the Geologic Evolution of the Pacific Northwest</u>, McGraw-Hill Book Co. (1972).
- 17. R. C. Newcomb, J. R. Strand, and F. J. Frank, <u>Geology and Groundwater Characteristics of the Hanford Reservation of the U.S. Atomic Energy Commission</u>, <u>Washington</u>, U.S. Geological Survey Prof. Paper 717 (1972).
- J. M. Parr, <u>Sieve Analysis Program</u>, Atlantic Richfield Hanford Company, Unpublished.
- 19. G. D. Webster, V. R. Barber, and C. Gustafson, <u>Channeled Scabland of Southeastern Washington A Road Log Via Spokane Coulee City Vantage Washtucna Lewiston Pullman</u>, Geological Society of America, Cordilleran Section, 72nd Annual Meeting, Field Guide No. 2 (1976).
- 20. C. K. Wentworth, <u>A Scale of Grade and Class Terms for Clastic Sediments</u>, Journal of Geology, V. 30, 377-392 (1922).

DISTRIBUTION

Energy Research and Development Administration - Richland Operations Office

0.J. Elgert Fed 700

J.A. Fernandez Fed 700

A.G. Lasila Fed 700

Battelle-Northwest

R.L. Brodzinski

J.R. Eliason

V.L. McGhan

S.J. Phillips

J. Raymond

R.W. Wallace

Atlantic Richfield Hanford Company

B.W. Anderson

J. Anderson

Η. Babad

G.E. Backman

L.D. Bockstanz

D.G. Bouse

D.J. Brown

L.E. Bruns

Burton, Jr. G.

R.A. Deju

F.R. Dornheim

G.T. Dukelow

P.G. Easley

K.R. Fecht

R.D. Fox

D.G. Harlow

W.M. Harty

0.F. Hill

H.F. Jensen

M.W. Legatski

B.J. McMurray

P.W. Metz

W.H. Price (2)

R.C. Roal

B.J. Saueressig


V.D. Schrag

H.P. Shaw

J.A. Teal

R.E. Van der Cook


R.L. Walser





<u>-1</u>-E27-75 (25 WEST) 30-03-03 SUGHTLY PEBBLY C-103 E27-76 (II'WEST) 30-03-05 SLIGHTLY PEBBLY VERY SLIGHTLY SILTY YERY COARSE TO COARSE SAND TO YERY COARSE SAND --H--SLIGHTLY SILTY COARSE TO MEDIUM SAUD CROSS SECTION SLIGHTLY PEBBLY VERY COARSE TO VERY COARSE TO COARSE SAND SLIGHTLY SILTY VERY COARSE TO COARSE TO COARSE SAND C-102 A-*X* STIGHTLY SILTY CO COARSE TO WEDIUM SAND SANO 5-101 C-101 -F-90 8 540 8 520 ¥ EXERCY RESEARCH 1 DEVELOPMENT ADMINISTRATION

NUMBER OF SECURITIES SITES

AND SECURITY SECURI ი







COARSE TO MEDIUM DAND E27-107 (42 EAST) PEBBLY VERY COARSE TO COARSE SAUD SLIGHTLY SHITY VERY COARSE TO CCARSE SAND TO VERY COARSE TO CCARSE SAND E27-109 (Z4 EAST) PEBBLY VERY COARSE TO COARSE SAND
COARSE TO COARSE
CARSE TO COARSE
SAND TO VERY FINE SAND STIGHTLY PEBBLY VERY COARSE TO COARSE SAND —H— CRoss VERY COARSE TO MEDIUM SAND SECTION E 27-10G (G'EAST) 30-11-11 Ċ.E Ш SUGHTLY SILTY COARSE TO MEDIUM SAID TO COARSE TO MEDIUM SAID TO ME Ш SILTY SANDY GRAVEL TO SUGHTLY SILTY WARSE TO COARSE SAND SANO TO COARSE TO MEDIUM SAND —G— E27-101 (35 EAST) PEBBLY YERY CCARSE SAND SLIGHTLY PEBBLY YERY COARSE TO COARSE TO TO PERBLY YERY COARSE TO /\/\ E27-103 (30'EAST) 30-10-09 -F-- ш<u>́</u> ğ 8 520 5 그를 돌 NONE O O

